

EXTENDED SCHUR FUNCTIONS AND BASES RELATED BY INVOLUTIONS

Spencer Daugherty · North Carolina State University · sdaughe@ncsu.edu

For a composition α , a **shin-tableau** T is a diagram of shape α filled with positive integers such that each row weakly increases from left to right and each column strictly increases from top to bottom.

The Extended Schur Functions [1]

For a composition α , the **extended Schur function** is defined by

$$\mathfrak{w}_\alpha^* = \sum_T x^T,$$

where the sum runs over shin-tableaux T of shape α . These functions form a basis of $QSym$, and \mathfrak{w}_λ^* is equal to the Schur function s_λ when λ is a partition.

$$\mathfrak{w}_{(2,3)}^* = x_1^2 x_2^3 + x_1^2 x_2^2 x_3 + x_1^2 x_2 x_3^2 + x_1 x_2^2 x_3^2 + x_1 x_2 x_3 x_4^2 + \dots$$

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & \\ \hline 2 & 2 & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & \\ \hline 2 & 3 & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & \\ \hline 2 & 4 & 4 \\ \hline \end{array} \quad \dots$$

A **standard** shin-tableau U of size n has entries $\{1, 2, \dots, n\}$ each appearing once.

$$Des_{\mathfrak{S}\mathfrak{w}}(U) = \{i : i + 1 \text{ is in a strictly lower row than } i\}$$

$$co_{\mathfrak{S}\mathfrak{w}}(U) = (i_1, i_2 - i_1, \dots, i_d - i_{d-1}, n - i_d) \text{ for } Des_{\mathfrak{S}\mathfrak{w}}(U) = \{i_1, \dots, i_d\}$$

For a composition α ,

$$\mathfrak{w}_\alpha^* = \sum_U F_{co_{\mathfrak{S}\mathfrak{w}}(U)},$$

where the sum runs over standard shin-tableaux U of shape α .

The **complement** of a composition α , denoted α^c , is the composition obtained from the complement of the set associated with α . The **reverse** of $(\alpha_1, \dots, \alpha_k)$, denoted α^r , is $(\alpha_k, \dots, \alpha_1)$. The **transpose** of α is defined by $\alpha^t = (\alpha^r)^c = (\alpha^c)^r$.

Involutions on $QSym$ [4]

For a composition α , define the following involutive automorphisms on $QSym$:

$$\psi(F_\alpha) = F_{\alpha^c} \quad \rho(F_\alpha) = F_{\alpha^r} \quad \omega(F_\alpha) = F_{\alpha^t}$$

Row-strict Extended Schur Functions [5]

The **row-strict extended Schur functions** are defined via *row-strict shin-tableaux* which have strictly increasing rows and weakly increasing columns. For a **standard** row-strict shin-tableau U , a **descent** $i \in Des_{\mathfrak{R}\mathfrak{w}}(U)$ is defined as an entry i such that $i + 1$ is in a weakly higher row.

The set of standard row-strict shin-tableaux is the same as the set of standard shin-tableaux, and for a standard (row-strict) shin-tableau U , we have $co_{\mathfrak{R}\mathfrak{w}}(U)^c = co_{\mathfrak{R}\mathfrak{w}}(U)$.

For a composition α ,

$$\psi(\mathfrak{w}_\alpha^*) = \mathfrak{R}\mathfrak{w}_\alpha^*$$

An Involution on the Schur Functions

The classical involution $\omega : Sym \rightarrow Sym$ maps the Schur basis to itself by $\omega(s_\lambda) = s_{\lambda'}$ where λ' is the conjugate of λ . Collectively, the involutions ψ , ρ , and ω on $QSym$ serve as an analogue to the classical ω . The extended Schur functions are part of what is essentially a system of four bases that is closed under the involutions ψ , ρ , and ω in $QSym$. **We introduce two new Schur-like bases of $QSym$** that, when paired with the extended Schur and row-strict extended Schur functions, complete this picture.

ψ and ω on $QSym$ restrict to ω on Sym , and ρ restricts to the identity map on Sym .

Flipped Extended Schur Functions

Let α be a composition and β a weak composition. A **flipped shin-tableau** of shape α and type β is a composition diagram α filled with positive integers that weakly decrease along the rows from left to right and strictly increase along the columns from top to bottom, where each positive integer i appears β_i times. A **standard** flipped tableau S of size n contains the entries $\{1, 2, \dots, n\}$ each exactly once.

$$Des_{\mathfrak{F}\mathfrak{w}}(S) = \{i : i + 1 \text{ is in a strictly lower row than } i\}$$

$$co_{\mathfrak{F}\mathfrak{w}}(S) = (i_1, i_2 - i_1, \dots, i_d - i_{d-1}, n - i_d) \text{ for } Des_{\mathfrak{F}\mathfrak{w}}(S) = \{i_1, \dots, i_d\}$$

For a composition α , the **flipped extended Schur function** is defined as

$$\mathfrak{F}\mathfrak{w}_\alpha^* = \sum_S F_{co_{\mathfrak{F}\mathfrak{w}}(S)},$$

where the sum runs over standard flipped shin-tableaux S of shape α .

There is a map *flip* between standard shin-tableaux U and standard flipped shin-tableaux S such that the descent composition of U is the reverse of the descent composition of $flip(U) = S$. First, flip U horizontally, then replace each entry i with $n - i$.

$$(2, 2, 1) \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 3 & 5 & \\ \hline 1 & 2 & 4 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 3 & 1 \\ \hline 5 & 4 \\ \hline \end{array} \quad (1, 2, 2)$$

Theorem (D. 2023)

For a composition α ,

$$\rho(\mathfrak{w}_\alpha^*) = \mathfrak{F}\mathfrak{w}_{\alpha^r}^* \text{ and } \omega(\mathfrak{R}\mathfrak{w}_\alpha^*) = \mathfrak{F}\mathfrak{w}_{\alpha^r}^*.$$

Backward Extended Schur Functions

Let α be a composition and β be a weak composition. A **backward shin-tableau** of shape α and type β is a composition diagram α filled with positive integers that strictly decrease along the rows from left to right and weakly increase along the columns from top to bottom, where each integer i appears β_i times. A **standard** backward tableau S of size n contains the entries $\{1, 2, \dots, n\}$ each exactly once.

$$Des_{\mathfrak{B}\mathfrak{w}}(S) = \{i : i + 1 \text{ is in a weakly higher row than } i\}$$

$$co_{\mathfrak{B}\mathfrak{w}}(S) = (i_1, i_2 - i_1, \dots, i_d - i_{d-1}, n - i_d) \text{ for } Des_{\mathfrak{B}\mathfrak{w}}(S) = \{i_1, \dots, i_d\}$$

For composition α , the **backward extended Schur function** is defined as

$$\mathfrak{B}\mathfrak{w}_\alpha^* = \sum_S F_{co_{\mathfrak{B}\mathfrak{w}}(S)},$$

where the sum runs over standard backward shin-tableaux S of shape α .

Theorem (D. 2023)

For a composition α ,

$$\omega(\mathfrak{w}_\alpha^*) = \mathfrak{B}\mathfrak{w}_{\alpha^r}^* \text{ and } \rho(\mathfrak{R}\mathfrak{w}_\alpha^*) = \mathfrak{B}\mathfrak{w}_{\alpha^r}^* \text{ and } \psi(\mathfrak{F}\mathfrak{w}_\alpha^*) = \mathfrak{B}\mathfrak{w}_\alpha^*.$$

Backward shin-tableaux are a row-strict version of flipped shin-tableaux. In fact, the set of standard row-strict tableaux and the set of standard backward tableaux are the same but for a tableau S , we will have $co_{\mathfrak{F}\mathfrak{w}}(S) = co_{\mathfrak{B}\mathfrak{w}}(S)^c$.

$$\mathfrak{w}_{(3,2)}^* = F_{(3,2)} + F_{(2,2,1)} + F_{(1,3,1)} + F_{(2,3)} + F_{(1,2,2)}$$

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array}$$

$$\mathfrak{R}\mathfrak{w}_{(3,2)}^* = F_{(1,1,2,1)} + F_{(1,2,2)} + F_{(2,1,2)} + F_{(1,2,1,1)} + F_{(2,2,1)}$$

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array}$$

$$\mathfrak{F}\mathfrak{w}_{(2,3)}^* = F_{(2,3)} + F_{(1,2,2)} + F_{(1,3,1)} + F_{(3,2)} + F_{(2,2,1)}$$

$$\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 5 & 4 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 3 & 1 & \\ \hline 5 & 4 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 4 & 1 & \\ \hline 5 & 3 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 3 & 2 & \\ \hline 5 & 4 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 4 & 2 & \\ \hline 5 & 3 & 1 \\ \hline \end{array}$$

$$\mathfrak{B}\mathfrak{w}_{(2,3)}^* = F_{(1,2,1,1)} + F_{(2,2,1)} + F_{(2,1,2)} + F_{(1,1,2,1)} + F_{(1,2,2)}$$

$$\begin{array}{|c|c|} \hline 2 & 1 \\ \hline 5 & 4 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 3 & 1 & \\ \hline 5 & 4 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 4 & 1 & \\ \hline 5 & 3 & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 3 & 2 & \\ \hline 5 & 4 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 4 & 2 & \\ \hline 5 & 3 & 1 \\ \hline \end{array}$$

Dual Bases in $NSym$

Each of our bases is dually paired with a variant of the *shin basis* of $NSym$. We introduce two of these: the **flipped shin functions** and **backward shin functions**. For each of these bases, we can obtain immediate properties by applying ψ , ρ , or ω to results on the shin functions. We also know the *commutative image* of a flipped or backward shin function. For a partition λ ,

$$\chi(\mathfrak{F}\mathfrak{w}_{\lambda^r}) = s_\lambda \text{ and } \chi(\mathfrak{B}\mathfrak{w}_{\lambda^r}) = s_\lambda,$$

and for a composition α that is not the reverse of a partition,

$$\chi(\mathfrak{F}\mathfrak{w}_\alpha) = \chi(\mathfrak{B}\mathfrak{w}_\alpha) = 0.$$

